

# *Molecular features of the basal-like breast cancer subtype based on BRCA1 mutation status*

**Aleix Prat, Cristina Cruz, Katherine A. Hoadley, Orland Díez, Charles M. Perou & Judith Balmaña**

**Breast Cancer Research and Treatment**

ISSN 0167-6806

Breast Cancer Res Treat  
DOI 10.1007/s10549-014-3056-x

Vol. 146 • No. 2 • July (I) 2014 • ISSN: 0167-6806

**ONLINE  
FIRST**

## **Breast Cancer**

*Research and Treatment*



 Springer  
the language of science

NO AUTHOR CHARGES FOR COLOR PRINTING

 Springer

**Your article is published under the Creative Commons Attribution Non-Commercial license which allows users to read, copy, distribute and make derivative works for noncommercial purposes from the material, as long as the author of the original work is cited. All commercial rights are exclusively held by Springer Science + Business Media. You may self-archive this article on your own website, an institutional repository or funder's repository and make it publicly available immediately.**

# Molecular features of the basal-like breast cancer subtype based on *BRCA1* mutation status

Aleix Prat · Cristina Cruz · Katherine A. Hoadley ·  
Orland Díez · Charles M. Perou · Judith Balmaña

Received: 23 January 2014 / Accepted: 5 July 2014  
© The Author(s) 2014. This article is published with open access at Springerlink.com

**Abstract** *BRCA1*-mutated breast cancer is associated with basal-like disease; however, it is currently unclear if the presence of a *BRCA1* mutation depicts a different entity within this subgroup. In this study, we compared the molecular features among basal-like tumors with and without *BRCA1* mutations. Fourteen patients with *BRCA1*-mutated (nine germline and five somatic) tumors and basal-like disease, and 79 patients with *BRCA1* non-mutated tumors and basal-like disease, were identified from the cancer genome atlas dataset. The following molecular data types were evaluated: global gene expression, selected protein and phospho-protein expression, global miRNA expression, global DNA methylation, total number of somatic mutations, *TP53* and *PIK3CA* somatic mutations, and global DNA copy-number aberrations. For intrinsic subtype identification, we used the PAM50 subtype predictor. Within the basal-like disease, we observed minor molecular differences in terms of gene, protein, and

miRNA expression, and DNA methylation variation, according to *BRCA1* status (either germline or somatic). However, there were significant differences according to average number of mutations and DNA copy-number aberrations, and four amplified regions (2q32.2, 3q29, 6p22.3, and 22q12.2), which are characteristic in high-grade serous ovarian carcinomas, were observed in both germline and somatic *BRCA1*-mutated breast tumors. These results suggest that minor, but potentially relevant, baseline molecular features exist among basal-like tumors according to *BRCA1* status. Additional studies are needed to better clarify if *BRCA1* genetic status is an independent prognostic feature, and more importantly, if *BRCA1* mutation status is a predictive biomarker of benefit from DNA-damaging agents among basal-like disease.

**Keywords** Basal-like · *BRCA1* · Intrinsic subtype · Breast cancer

## Abbreviations

TN Triple-negative  
*BRCA1* Breast cancer 1 early onset

Aleix Prat and Cristina Cruz have contributed equally to this work.

**Electronic supplementary material** The online version of this article (doi:10.1007/s10549-014-3056-x) contains supplementary material, which is available to authorized users.

A. Prat (✉)  
Translational Genomics Group, Vall d'Hebron Institute of  
Oncology (VHIO), Pg Vall d'Hebron, 119-129, 08035 Barcelona,  
Spain  
e-mail: aprat@vhio.net

C. Cruz · J. Balmaña  
High Risk and Cancer Prevention Group, Vall d'Hebron Institute  
of Oncology (VHIO), 08035 Barcelona, Spain  
e-mail: ccruz@vhio.net

J. Balmaña  
e-mail: jbalmana@vhio.net

K. A. Hoadley · C. M. Perou  
Lineberger Comprehensive Cancer Center, University of North  
Carolina at Chapel Hill, Chapel Hill, NC 27514, USA  
e-mail: hoadley@med.unc.edu

C. M. Perou  
e-mail: cperou@med.unc.edu

O. Díez  
Oncogenetics Group, Hospital Universitari de la Vall d'Hebron,  
Vall d'Hebron Institute of Oncology (VHIO), 08035 Barcelona,  
Spain  
e-mail: odiez@vhebron.net

ID4	Inhibitor of DNA binding 4, dominant negative helix-loop-helix protein
PIK3CA	Phosphatidylinositol-4, 5-bisphosphate 3-kinase, catalytic subunit alpha

methylation variation: “*BRCA.methylation.27 k.450 k.txt.*” For microarray DNA copy-number aberration data: “*brca\_scna\_all\_thresholded.by\_genes.txt.*” For intrinsic subtype identification, we used the PAM50 subtype calls as provided in the TCGA portal.

#### Independent dataset

We evaluated an independent and publicly available microarray-based gene expression dataset (GSE40115) that includes breast tumors from 32 patients with basal-like disease (20 with *BRCA1* germline mutations and 12 with sporadic tumors [i.e. unknown *BRCA1* status]). The file “*GSE40115-GPL15931\_series\_matrix.txt*” with the normalized log<sub>2</sub> ratios (Cy5 sample/Cy3 control) of probes was used. Probes mapping to the same gene (Entrez ID as defined by the manufacturer) were averaged to generate independent expression estimates.

#### Seven-TN subtype classification

To identify the 7-TN subtypes described by Lehmann et al. [16], (i.e., basal one, basal two, immunomodulatory, luminal androgen receptor, mesenchymal, mesenchymal stem cell, and unstable), we submitted the raw gene expression data of each individual dataset of basal-like disease to the TNBC type online predictor (<http://cbc.mc.vanderbilt.edu/tnbc/>) [17].

#### Statistical analysis

All multiple-testing comparisons were done using an unpaired two-class significance analysis of microarrays (SAM, <http://www-stat.stanford.edu/~tibs/SAM/>). The mutation rates of *TP53* and *PIK3CA* genes between two groups, the 7-TN subtype distribution between *BRCA1*-mutated and non-mutated basal-like tumors, and the amplification rates of *ID4* between two groups, were compared using the Chi square and Fisher’s exact tests. The total number of somatic mutations between two groups was compared using a Student’s *t* test. All statistical computations were performed in R v.2.15.1 (<http://cran.r-project.org>).

## Results and discussion

From TCGA breast cancer dataset, we identified 12 tumor samples with *BRCA1* germline mutations (all classified as deleterious), seven tumor samples with somatic *BRCA1* mutations, and one tumor sample with both *BRCA1* germline and somatic mutations (Supplemental Material). As expected, 70 % of *BRCA1* mutated tumors were of the basal-like intrinsic subtype (nine germline and five

## Introduction


Studies based on gene expression data have identified and characterized four main intrinsic subtypes of breast cancer (luminal A, luminal B, HER2-enriched, and basal-like) [1, 2]. Among them, the basal-like subtype is associated with young age, *BRCA1* germline and somatic mutations [1, 3, 4] and an overall poor prognosis despite that a subgroup of patients with these tumors has an excellent outcome when treated with chemotherapy [5]. In the clinical setting, basal-like tumors are usually identified by the lack of expression of hormone receptors by immunohistochemistry (IHC) and lack of overexpression of HER2 by IHC and/or FISH (the so called triple-negative [TN] status) [1, 2, 6]. Although the TN definition enriches for basal-like disease, considerable discordance exists [2, 6].

*BRCA1* mutations and other associated molecular traits might confer sensitivity to specific therapeutic agents [7–10]. Nevertheless, it is unclear how different, from a biological perspective, *BRCA1*-mutated basal-like tumors are from *BRCA1* non-mutated basal-like tumors, and whether *BRCA1* mutation is an independent prognostic and/or predictive biomarker when the intrinsic subtype is taken into account [11–15]. This line of thought directed us to formulate the question of how much the biology of basal-like tumors with *BRCA1* mutations differs from the biology of basal-like tumors without *BRCA1* mutations. To address this question, we interrogated The Cancer Genome Atlas (TCGA) breast cancer project which provides various types of molecular data coming from DNA, RNAs, and proteins [1].

## Methods

### The Cancer Genome Atlas dataset

In this study, we evaluated TCGA breast cancer dataset and all data were obtained from the TCGA breast cancer online portal ([https://tcga-data.nci.nih.gov/docs/publications/brca\\_2012/](https://tcga-data.nci.nih.gov/docs/publications/brca_2012/)). The following files were used. For microarray gene expression data: “*BRCA.exp.547.med.txt.*” For reverse-phase protein array (RPPA) expression data: “*rppaData-403Samp-171Ab-Trimmed.txt.*” For sequencing miRNA expression: “*BRCA.780.mimat.txt.*” For microarray DNA



**Fig. 1** Intrinsic profile of *BRCA1*-mutated breast tumors. Hierarchical clustering of 509 breast samples of the cancer genome atlas (TCGA) project using the ~1,900 intrinsic gene list [30]. PAM50 intrinsic subtype calls [30] and *BRCA1* mutation status is shown below the array tree


**Table 1** Significant molecular differences between basal-like *BRCA1*-mutated tumors ( $n = 14$ ) and basal-like *BRCA1* non-mutated tumors ( $n = 79$ )

Total biomarkers evaluated	Type of evaluation	Comparison (more expressed or amplified)	Significant biomarkers identified (FDR = 0 %)	Percentage of altered biomarkers (%)
17,786 (unique genes)	Expression	<i>BRCA1</i> MUT <i>BRCA1</i> WT	0 0	0
171 (unique proteins or phospho-proteins by RPPA)	Expression	<i>BRCA1</i> MUT <i>BRCA1</i> WT	0 1	0.6
1,222 (mature/star miRNA strands)	Expression	<i>BRCA1</i> MUT <i>BRCA1</i> WT	3 0	0.2
530 (unique genes)	Methylation	<i>BRCA1</i> MUT <i>BRCA1</i> WT	0 6	1.1
19,613 (unique genes)	DNA amplification	<i>BRCA1</i> MUT <i>BRCA1</i> WT	250 0	1.3

RPPA reverse-phase protein arrays, FDR false discovery rate, *BRCA1*WT *BRCA1* wild-type, *BRCA1*MUT *BRCA1* mutated

somatic), but luminal A (two germline, one germline/somatic, and one somatic), luminal B (one germline), and HER2-enriched (one somatic) tumors were also identified (Fig. 1). Similarly, 66.7 % of *BRCA1* mutated tumors were TN.

Within basal-like disease, we observed minor molecular differences (0–1.1 %) in terms of gene expression, protein expression, miRNA expression, and DNA methylation variation according to *BRCA1* status (Table 1 and Supplemental Material). Indeed, no genes among 17,876 genes were found differentially expressed between basal-like *BRCA1*-mutated tumors versus basal-like *BRCA1* non-mutated tumors (Table 1), including the *BRCA1* mRNA transcript (Fig. 2). Similar results were observed when only the tumors with *BRCA1* germline mutations were taken into consideration (Supplemental Material). Concordant with this result, analysis of microarray gene expression data of an independent dataset of 32 tumors with basal-like disease (20 with a *BRCA1* germline mutation and 12 with sporadic tumors) revealed only 0.03 % differentially expressed genes (6 of 21,848, false discovery rate



**Fig. 2** Relative *BRCA1* gene expression in basal-like disease based on *BRCA1* mutational status. Data have been obtained from the TCGA breast cancer project. The *BRCA1* gene expression has been median centered across all breast cancer samples with DNA-seq data (i.e., basal-like and not basal-like). The  $p$ -value was calculated by comparing gene expression means across the three groups. In red color, breast samples with  $\geq 2$ -fold decrease in *BRCA1* expression compared to its median expression in breast cancer are shown

**Table 2** DNA regions found significantly more amplified in basal-like *BRCA1*-mutated tumors ( $n = 14$ ) compared to basal-like *BRCA1* non-mutated tumors ( $n = 74$ )

Basal-like <i>BRCA1</i> mutated	HGSOC	Genes
6p22.3	6p22.3	FAM65B, TDP2, ACOT13, ALDH5A1, GPLD1, KIAA0319, MRS2, C6orf62, GMNN, DCDC2, CMAHP, KAAG1, KIF13A, DEK, NRSN1, E2F3, MBOAT1, RNF144B, CDKAL1, KDM1B, NHLRC1, TPMT, ID4, HDGFL1, PRL, LINC00340, SOX4, CAP2, FAM8A1, NUP153, RBM24, MYLIP, GMPR, ATXN1, DTNBP1, JARID2
3q29	3q29	FYTTD1, KIAA0226, DLG1, BDH1, LOC220729, CEP19, LOC152217, MFI2, NCBP2, PAK2, PIGX, PIGZ, SENP5, ACAP2, ANKRD18DP, FAM157A, LMLN, IQCG, LRCH3, C3orf43, FBXO45, LRRC33, RNF168, UBXN7, WDR53, APOD, MUC20, MUC4, OSTalpha, PCYT1A, PPP1R2, SDHAP1, SDHAP2, TCTEX1D2, TFRC, TM4SF19, TNK2, ZDHHC19, XXYL1, FAM43A, LSG1, TMEM44, RPL35A, ATP13A3, ATP13A4, ATP13A5, CPN2, GP5, HES1, HRASLS, LOC100128023, LOC100131551, LRRC15, MB21D2, MGC2889, OPA1
2q32.2	2q32.2	COL3A1, COL5A2, DIRC1, NAB1, TMEM194B, C2orf88, GLS, HIBCH, INPP1, MFSD6, MSTN, STAT4, SLC40A1, WDR75, ORMDL1, OSGEPL1, PMS1, ANKAR, ASNSD1, STAT1
22q12.2	22q12.2	APIB1, ASCC2, CABP7, CCDC157, DEPDC5, DRG1, DUSP18, EIF4ENIF1, EMID1, EWSR1, GAL3ST1, GAS2L1, GATSL3, HORMAD2, INPP5 J, LIF, LIMK2, MORC2, MORC2-AS1, MTFP1, MTMR3, NEFH, NF2, NIPSNAP1, OSBP2, OSM, PATZ1, PES1, PIK3IP1, PISD, PLA2G3, PRR14L, RASL10A, RFPL1, RFPL1-AS1, RHBDD3, RNF185, RNF215, SDC4P, SEC14L2, SEC14L3, SEC14L4, SELM, SF3A1, SFI1, SLC35E4, SMTN, SNORD125, TBC1D10A, TCN2, THOC5, TUG1, UQCR10, ZMAT5
10q25.3	–	TRUB1, CASP7, ATRNL1, FAM160B1, PDZD8, SLC18A2, C10orf96, C10orf81, DCLRE1A, HABP2, NHLRC2, NRAP, KCNK18, KIAA1598, VAX1, GFRA1, PNLIP, PNLIPRP1, PNLIPRP2, PNLIPRP3, C10orf82, HSPA12A, ADRB1, AFAP1L2, C10orf118, TDRD1, VWA2, ABLIM1
10q26.11	–	PRLHR, FAM204A, BAG3, INPP5F, TIAL1, C10orf46, MCMBP, SEC23IP, CASC2, EMX2, EMX2OS, RAB11FIP2, EIF3A, FAM45A, GRK5, NANOS1, PRDX3, RGS10, SFXN4, SNORA19
22q11.22	–	GGTLC2, GNAZ, LOC648691, LOC96610, POM121L1P, PPM1F, PRAME, RAB36, RTDR1, TOP3B, VPREB1, ZNF280A, ZNF280B
22q11.23	–	ADORA2A, BCR, BCRP3, C22orf13, C22orf15, C22orf43, C22orf45, CABIN1, CHCHD10, CRYBB2, CRYBB3, DDT, DDTL, DERL3, FAM211B, GGT1, GGT5, GSTT1, GSTT2, GSTTP1, GSTTP2, GUSBP11, IGLL1, IGLL3P, KIAA1671, LOC391322, LRP5L, MIF, MMP11, PIWIL3, POM121L10P, POM121L9P, RGL4, SGSM1, SLC2A11, SMARCB1, SNRPD3, SPECC1L, SUSD2, TMEM211, TOP1P2, UPB1, VPREB3, ZDHHC8P1, ZNF70
22q12.1	–	ADRBK2, ASPHD2, C22orf31, CCDC117, CHEK2, CRYBA4, CRYBB1, HPS4, HSCB, KREMEN1, MIAT, MN1, MYO18B, PITPNB, SEZ6L, SRRD, TFIP11, TPST2, TTC28, TTC28-AS1, XBP1, ZNRF3
22q12.3	–	C22orf24, C22orf42, SLC5A1, YWHAH, BPIFC, C22orf28, FBXO7, RFPL2, RFPL3, RFPL3-AS1, SLC5A4, SYN3, APOL5, APOL6, HMOX1, MB, MCM5, RASD2, TOM1, TIMP3, CACNG2, IFT27, PVALB, NCF4, C1QTNF6, C22orf33, CSF2RB, IL2RB, KCTD17, MPST, TMPRSS6, TST, ISX, HMGXB4, LARGE, APOL3, RBFOX2, EIF3D, FOXRED2, TXN2, APOL1, MYH9, APOL2, APOL4
2q32.1	–	ZSWIM2, ZNF804A, FAM171B, ITGAV, GULP1, CALCRL, TFPI, ZC3H15, DNAJC10, DUSP19, NUP35, FRZB, NCKAP1, PDE1A
2q33.2	–	CTLA4, ICOS, CD28, RAPH1, FAM117B, ICA1L, ABI2, ALS2CR8, WDR12, CYP20A1, NBEAL1
3q28	–	CCDC50, FGF12, OSTN, PYDC2, UTS2D, CLDN1, CLDN16, GMNC, IL1RAP, LEPREL1, SNAR-I, TMEM207, TP63, TPRG1
6p21.31	–	NUDT3, C6orf1, HMGA1, BAK1, GGNBP1, LINC00336, ANKS1A, C6orf126, C6orf127, C6orf81, CLPS, FKBP5, GRM4, LHFPL5, LOC285847, SCUBE3, SNRPC, SRPK1, TAF11, TCP11, UHRF1BP1, SLC26A8, C6orf125, IP6K3, ITPR3, LEMD2, MLN, RPL10A, TEAD3, TULP1, ZNF76, C6orf106, PACSIN1, RPS10, SPDEF, BRPF3, C6orf222, MAPK13, MAPK14, PNPLA1, DEF6, FANCE, PPARD, ETV7, PXT1, KCTD20, SRSF3, STK38

HGSOC high-grade serous ovarian carcinoma

[FDR] = 0 %) between the two groups [18] (Supplemental Material). In addition, we did not identify significant differences in the proportion of the recently reported 7-TN subtype classification proposed by Lehmann and colleagues [16], between basal-like tumors with and without *BRCA1* mutations (Supplemental Material). Interestingly, two clear groups within the basal-like *BRCA1* wild-type

disease were identified based on *BRCA1* mRNA expression-only (i.e., high and low) (Fig. 2).

In terms of DNA copy-number aberrations, we identified 250 genes (representing 14 different DNA regions and 1.3 % of all genes evaluated) showing higher amplification rates in basal-like *BRCA1*-mutated tumors compared to basal-like *BRCA1* wild-type tumors (Table 2). Among

them, we identified four regions (2q32.2, 3q29, 6p22.3, and 22q11.2) that have been previously shown to be amplified and characteristic of high-grade serous ovarian carcinomas [19]. Interestingly, region 6p22.3 contains *ID4*, a gene long known to be a marker of basal-like breast cancers [20], and known to code for a DNA-binding protein that negatively regulates *BRCA1* expression in breast and ovarian cancers [21]. This gene was found amplified (i.e. low or high gains) in 78.6 % (11/14) of basal-like *BRCA1* mutated tumors versus 35.1 % (26/74) of basal-like *BRCA1* wild-type tumors ( $p = 0.008$ , Fisher's exact test). Similar results were observed when the *BRCA1* somatic mutations were excluded (Supplemental Material). The biological role of *ID4* amplification in *BRCA1* mutated breast cancer is currently unknown, and we could hypothesize that *ID4* might inhibit residual function of mutant *BRCA1*.

In terms of somatic gene mutations, basal-like *BRCA1* mutated tumors showed higher average number of mutations than basal-like *BRCA1* wild-type tumors (122.6 vs. 80.3,  $p = 0.004$ , Student's *t* test). Regarding the distribution of *TP53* and *PIK3CA* somatic mutations according to *BRCA1* status, *TP53* mutations were found in 100 % (14/14) of basal-like *BRCA1* mutated versus 75.9 % (60/79) of basal-like *BRCA1* wild-type tumors ( $p = 0.065$ , Fisher's exact test). Finally, *PIK3CA* mutations were found in 0 % (0/14) of basal-like *BRCA1* mutated tumors versus 10.1 % (8/79) of basal-like *BRCA1* wild-type tumors ( $p = 0.602$ ).

In our analysis, most of the unique molecular features of basal-like *BRCA1* mutated tumors were found at the DNA level (i.e. amplifications and mutation rates). Indeed, basal-like *BRCA1* mutated tumors showed higher amplification rates at 14 different chromosomal regions and higher number of somatic mutations, including *TP53*, compared to basal-like *BRCA1* wild-type tumors. However, no significant differences in protein expression were found when comparing basal-like *BRCA1* mutated and *BRCA1* wild-type tumors. These results suggest that the genomic instability induced by *BRCA1* loss [22] does not translate into a recognizable phenotype at the RNA and protein level. The potential explanation of these findings is currently unknown. Nonetheless, the fact that 4 out of 14 (28.5 %) amplified DNA regions were found to be characteristic regions of high-grade serous ovarian carcinomas suggests that, among basal-like breast tumors, those with a *BRCA1* mutation are more similar to ovarian carcinoma at the genetic level.

In our analysis, the absence of recognizable prominent differences in molecular alterations based on *BRCA1* mutation status would be in line with previous clinical data suggesting that *BRCA1* status *per se* might not play a major role in conferring a distinct prognosis within basal-like disease. Results from three retrospective studies that have

evaluated the prognostic role of *BRCA1/2* mutations (mostly *BRCA1*) in TN breast cancer support this hypothesis [13–15]. In Bayraktar et al. [13], *BRCA1/2* status was not found to be prognostic in 227 women with early TN breast cancer referred to genetic counseling. Similar results were observed in a cohort of 195 patients with metastatic breast cancer, where the independent prognostic value of *BRCA1* in univariate analyses was lost when TN status and other clinical-pathological variables were taken into account [14]. More recently, Huzarski et al. [15] evaluated the association of germline *BRCA1* mutation status with 10 year overall survival in 3,350 polish women with a diagnosis of breast cancer. The authors observed that *BRCA1* mutation status was significantly associated with worse outcome when standard clinical-pathological variables were taken into account [15]. However, among patients with TN breast cancer, *BRCA1* status was not associated with worse outcome [15].

The role of the *BRCA1* mutation status as a predictive factor of treatment response among TN breast cancer is also under study. On the one hand, two retrospective studies have evaluated the ability of *BRCA1* mutation status to predict response to multi-agent chemotherapy [11, 12]. In the first study, Arun and colleagues showed no significant differences in terms of pathological complete response rates after neoadjuvant chemotherapy (mostly anthracycline/taxane-based) among 75 patients with TN breast cancer in relation to their *BRCA1* status [11]. In the second study, Gonzalez-Angulo et al. [12] observed a better outcome in *BRCA1/2* mutated TN breast cancer compared to *BRCA1/2* non-mutated TN breast cancers after treatment with adjuvant anthracycline/taxane-based chemotherapy. On the other hand, two recent prospective clinical trials (GeparSixto [23] and CALGB40603 [24]) have demonstrated the value of adding carboplatin, a DNA-damaging agent, to standard neoadjuvant anthracycline/taxane-based chemotherapy in 769 patients with newly diagnosed TN breast cancer, regardless of their *BRCA1* mutational status.

Previous retrospective studies have suggested that *BRCA1* mutated tumors might substantially benefit from platinum [9, 25]. In fact, in the GeparSixto TN trial [23, 26], recent data reported higher pCR rates in *BRCA1/2*-mutated patients compared to *BRCA1/2* non-mutated patients. Nevertheless, data on the intrinsic subtype of the TN wild-type tumors in this clinical trial have not been reported yet and it might be interesting to analyze whether the basal-like benefits the most. Supporting the hypothesis that basal-like *BRCA1* non-mutated breast cancers might also benefit to some extent from DNA-damaging agents, several studies have identified *BRCA1* mutation-unrelated mechanisms of platinum sensitivity in TN *BRCA1* wild-type breast cancer such as the p63/p73 network, telomeric

allelic imbalance, and homologous recombination deficiency [27–29].

## Conclusions

In this study, we compared DNA, RNA, and protein data among basal-like tumors with and without *BRCA1* mutations and observed that minor molecular features exist. The clinical relevance of these differences is unknown and further validation in larger and prospective cohorts is warranted. Biomarker analyses are needed to clarify if *BRCA1* status is an independent prognostic feature and/or a predictive biomarker of benefit from DNA-damaging agents beyond the basal-like phenotype.

**Acknowledgments** Beca Roche en Onco-Hematología 2012

**Conflict of interest** C.M.P is an equity stock holder, and Board of Director Member, of BioClassifier LLC and University Genomics. C.M.P is also listed an inventor on a patent application on the PAM50 molecular assay. Uncompensated advisory role of A.P. for Nanostring Technologies.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

- TCGA (2012) Comprehensive molecular portraits of human breast tumours. *Nature* 490:61–70
- Prat A, Adamo B, Cheang MCU, Anders CK, Carey LA, Perou CM (2013) Molecular characterization of basal-like and non-basal-like triple-negative breast cancer. *Oncologist* 18(2):123–133. doi:10.1634/theoncologist.2012-0397
- Foulkes WD, Stefansson IM, Chappuis PO, Begin LR, Goffin JR, Wong N, Trudel M, Akslen LA (2003) Germline *BRCA1* mutations and a basal epithelial phenotype in breast cancer. *J Natl Cancer Inst* 95:1482–1485
- Lee E, McKean-Cowdin R, Ma H, Spicer DV, Van Den Berg D, Bernstein L, Ursin G (2011) Characteristics of triple-negative breast cancer in patients with a *BRCA1* mutation: results from a population-based study of young women. *J Clin Oncol* 29(33):4373–4380. doi:10.1200/jco.2010.33.6446
- Carey LA, Dees EC, Sawyer L, Gatti L, Moore DT, Collichio F, Ollila DW, Sartor CI, Graham ML, Perou CM (2007) The triple negative paradox: primary tumor chemosensitivity of breast cancer subtypes. *Clin Cancer Res* 13(8):2329–2334. doi:10.1158/1078-0432.ccr-06-1109
- Goldhirsch A, Winer EP, Coates AS, Gelber RD, Piccart-Gebhart M, Thürlimann B, Senn H-J (2013) Personalizing the treatment of women with early breast cancer: highlights of the St Gallen international expert consensus on the primary therapy of early breast cancer 2013. *Ann Oncol*. doi:10.1093/annonc/mdt303
- Fong PC, Boss DS, Yap TA, Tutt A, Wu P, Mergui-Roelvink M, Mortimer P, Swaisland H, Lau A, O'Connor MJ, Ashworth A, Carmichael J, Kaye SB, Schellens JHM, de Bono JS (2009) Inhibition of poly(ADP-ribose) polymerase in tumors from *BRCA* mutation carriers. *N Engl J Med* 361(2):123–134. doi:10.1056/NEJMoa0900212
- Carey LA (2010) Targeted chemotherapy? platinum in *BRCA1*-dysfunctional breast cancer. *J Clin Oncol* 28(3):361–363. doi:10.1200/jco.2009.24.0838
- Byrski T, Gronwald J, Huzarski T, Grzybowska E, Budryk M, Stawicka M, Mierzwa T, Szwiec M, Wiśniowski R, Siolek M, Dent R, Lubinski J, Narod S (2010) Pathologic complete response rates in young women with *BRCA1*-positive breast cancers after neoadjuvant chemotherapy. *J Clin Oncol* 28(3):375–379. doi:10.1200/jco.2008.20.7019
- Balmaña J, Domchek SM, Tutt A, Garber JE (2011) Stumbling blocks on the path to personalized medicine in breast cancer: the case of PARP inhibitors for *BRCA1/2*-associated cancers. *Cancer Discov*. doi:10.1158/2159-8274.cd-11-0048
- Arun B, Bayraktar S, Liu DD, Gutierrez Barrera AM, Atchley D, Pusztaï L, Litton JK, Valero V, Meric-Bernstam F, Hortobagyi GN, Albarracín C (2011) Response to neoadjuvant systemic therapy for breast cancer in *BRCA* mutation carriers and non-carriers: a single-institution experience. *J Clin Oncol* 29(28):3739–3746. doi:10.1200/jco.2011.35.2682
- Gonzalez-Angulo AM, Timms KM, Liu S, Chen H, Litton JK, Potter J, Lanchbury JS, Stemke-Hale K, Hennessy BT, Arun BK, Hortobagyi GN, Do K-A, Mills GB, Meric-Bernstam F (2011) Incidence and outcome of *BRCA* mutations in unselected patients with triple receptor-negative breast cancer. *Clin Cancer Res* 17(5):1082–1089. doi:10.1158/1078-0432.ccr-10-2560
- Bayraktar S, Gutierrez-Barrera A, Liu D, Tasbas T, Akar U, Litton J, Lin E, Albarracín C, Meric-Bernstam F, Gonzalez-Angulo A, Hortobagyi G, Arun B (2011) Outcome of triple-negative breast cancer in patients with or without deleterious *BRCA* mutations. *Breast Cancer Res Treat* 130(1):145–153. doi:10.1007/s10549-011-1711-z
- Bayraktar S, Gutierrez-Barrera AM, Lin H, Elsayegh N, Tasbas T, Litton JK, Ibrahim NK, Morrow PK, Green M, Valero V, Booser DJ, Hortobagyi GN, Arun BK (2013) Outcome of metastatic breast cancer in selected women with or without deleterious *BRCA* mutations. *Clin Exp Metastasis* 30(5):631–642. doi:10.1007/s10585-013-9567-8
- Huzarski T, Byrski T, Gronwald J, Górski B, Domagała P, Cybulski C, Oszurek O, Szwiec M, Gugala K, Stawicka M, Morawiec Z, Mierzwa T, Janiszewska H, Kilar E, Marczyk E, Kozak-Klonowska B, Siolek M, Surdyka D, Wiśniowski R, Posmyk M, Sun P, Lubiński J, Narod SA (2013) Ten-year survival in patients with *BRCA1*-negative and *BRCA1*-positive breast cancer. *J Clin Oncol*. doi:10.1200/jco.2012.45.3571
- Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, Pietenpol JA (2011) Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. *J Clin Invest* 121(7):2750–2767. doi:10.1172/jci45014
- Chen X, Li J, Gray WH, Lehmann BD, Bauer JA, Shyr Y, Pietenpol JA (2012) TNBCtype: a subtyping tool for triple-negative breast cancer. *Cancer Inform* 11 (3284-CIN-TNBCtype:-A-Subtyping-Tool-for-Triple-Negative-Breast-Cancer.pdf):147–156. doi:10.4137/cin.s9983
- Larsen MJ, Kruse TA, Tan Q, Lænkholm A-V, Bak M, Lykkesfeldt AE, Sørensen KP, TvO Hansen, Ejlersen B, Gerdes A-M, Thomassen M (2013) Classifications within molecular subtypes enables identification of *BRCA1/BRCA2* mutation



- carriers by rna tumor profiling. PLoS ONE 8(5):e64268. doi:[10.1371/journal.pone.0064268](https://doi.org/10.1371/journal.pone.0064268)
19. TCGA (2011) Integrated genomic analyses of ovarian carcinoma. *Nature* 474:609–615
  20. Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, Pollack JR, Ross DT, Johnsen H, Akslen LA, Fluge O, Pergamenschikov A, Williams C, Zhu SX, Lonning PE, Borresen-Dale A-L, Brown PO, Botstein D (2000) Molecular portraits of human breast tumours. *Nature* 406(6797):747–752
  21. Beger C, Pierce LN, Krüger M, Marcusson EG, Robbins JM, Welch P, Welch PJ, Welte K, King M-C, Barber JR, Wong-Staal F (2001) Identification of Id4 as a regulator of *BRCA1* expression by using a ribozyme-library-based inverse genomics approach. *Proc Natl Acad Sci* 98(1):130–135. doi:[10.1073/pnas.98.1.130](https://doi.org/10.1073/pnas.98.1.130)
  22. Deng C-X (2006) *BRCA1*: cell cycle checkpoint, genetic instability, DNA damage response and cancer evolution. *Nucleic Acids Res* 34(5):1416–1426. doi:[10.1093/nar/gkl010](https://doi.org/10.1093/nar/gkl010)
  23. von Minckwitz G, Schneeweiss A, Loibl S, Salat C, Denkert C, Rezai M, Blohmer JU, Jackisch C, Paepke S, Gerber B, Zahm DM, Kümmel S, Eidtmann H, Klare P, Huober J, Costa S, Tesch H, Hanusch C, Hilfrich J, Khandan F, Fasching PA, Sinn BV, Engels K, Mehta K, Nekljudova V, Untch M (2014) Neoadjuvant carboplatin in patients with triple-negative and HER2-positive early breast cancer (GeparSixto; GBG 66): a randomised phase 2 trial. *Lancet Oncol* 15(7):747–756. doi:[10.1016/S1470-2045\(14\)70160-3](https://doi.org/10.1016/S1470-2045(14)70160-3)
  24. Sikov W, Berry D, Perou C, Singh B, Cirincione C, Tolaney S, Kuzma C, Pluard T, Somlo G, Port E, Golshan M, Bellon J, Collyar D, Hahn O, Carey L, Hudis C, Winer E (2013) Impact of the addition of carboplatin (Cb) and/or bevacizumab (B) to neoadjuvant weekly paclitaxel (P) followed by dose-dense AC on pathologic complete response (pCR) rates in triple-negative breast cancer (TNBC): CALGB 40603 (Alliance). San Antonio Breast Cancer Symposium S5-01
  25. Byrski T, Dent R, Blecharz P, Foszczynska-Kloda M, Gronwald J, Huzarski T, Cybulski C, Marczyk E, Chrzan R, Eisen A, Lubinski J, Narod S (2012) Results of a phase II open-label, non-randomized trial of cisplatin chemotherapy in patients with *BRCA1*-positive metastatic breast cancer. *Breast Cancer Res* 14(4):R110
  26. Von Minckwitz G, Hahnen E, Fasching P, Hauke J, Schneeweiss A, Salat C, Rezai M, Blohmer J, Zahm D, Jackisch C, Gerber B, Klare P, Kümmel S, Eidtmann H, Paepke S, Nekljudova V, Loibl S, Untch M, Schmutzler R, Groups aGaa-BS Pathological complete response (pCR) rates after carboplatin-containing neoadjuvant chemotherapy in patients with germline *BRCA* (gBRCA) mutation and triple negative breast cancer (TNBC)—Results from GeparSixto. *Proc Am Soc Clin Oncol*: a1005, 2014
  27. Leong C-O, Vidnovic N, DeYoung MP, Sgroi D, Ellisen LW (2007) The p63/p73 network mediates chemosensitivity to cisplatin in a biologically defined subset of primary breast cancers. *J Clin Investig* 117(5):1370–1380. doi:[10.1172/JCI30866](https://doi.org/10.1172/JCI30866)
  28. Birkbak NJ, Wang ZC, Kim J-Y, Eklund AC, Li Q, Tian R, Bowman-Colin C, Li Y, Greene-Colozzi A, Iglehart JD, Tung N, Ryan PD, Garber JE, Silver DP, Szallasi Z, Richardson AL (2012) Telomeric allelic imbalance indicates defective DNA repair and sensitivity to DNA-damaging agents. *Cancer Discov* 2(4):366–375. doi:[10.1158/2159-8290.cd-11-0206](https://doi.org/10.1158/2159-8290.cd-11-0206)
  29. Isakoff S, He L, Mayer E, Goss P, Traina T, Carey L, Krag K, Liu M, Rugo H, Stearns V, Come S, Ryan P, Finkelstein D, Hartman A, Garber J, Timms K, Winer E, Ellisen L Identification of biomarkers to predict response to single-agent platinum chemotherapy in metastatic triple-negative breast cancer (mTNBC): Correlative studies from TBCRC009. *Proc Am Soc Clin Oncol*: a1020, 2014
  30. Parker JS, Mullins M, Cheang MCU, Leung S, Voduc D, Vickery T, Davies S, Fauron C, He X, Hu Z, Quackenbush JF, Stijleman IJ, Palazzo J, Marron JS, Nobel AB, Mardis E, Nielsen TO, Ellis MJ, Perou CM, Bernard PS (2009) Supervised risk predictor of breast cancer based on intrinsic subtypes. *J Clin Oncol* 27(8):1160–1167. doi:[10.1200/jco.2008.18.1370](https://doi.org/10.1200/jco.2008.18.1370)