





2







5



	•
Aimee Faso, Plawno, CPROCP, is a Clinical Pharmacist Practitioner (CPP) at the North Carolina Reacight	
Almee Faso, Pharmo, CPR BCOP, is a Clinical Pharmacist Practitioner (CPP) at the North Carolina Basnight Cancer Hospital (NOSEH) who works with the breast oncology team. After completing her Master's degree in physiological tools upschology at Choi University, she worked in the pharmaceutical industry for 2 years. She then consider the control of the Completing Programmer of the Completing	
oncology team. After completing her Master's degree in physiological polychology at Olin University, she worked in the pharmaceutical industry for 2 years. She then in the pharmaceutical industry for 2 years. She then in Gainesville, F. I followed by a pharmacy praviate residency at Dartmouth Hitchcock Medical Center. Dr. Faso started working at the NCBCH in 2008 and in 2014 joined the outpatient breast oncology team. She where she teaches breast cancer treatment to pharmacy students and is a preceptor to pharmacy students and pharmacy residents. She is also a Clinical Assistant Professor in the School of Nursing where she teaches	_
Almee Faso, Paumit, ROR OP Paumit, ROR OP Dr. Faso started working at the NCBCH in 2008 and in 2014 joined the outpatient breast oncology team. She where she teaches breast cancer treatment to pharmacy students and is a preceptor to pharmacy students and pharmacy residents. She is also a Olinical Assistant Professor in the School of Nursing where she teaches oncology pharmacology in the nurse practitioner program.	
students and is a preceptor to pharmacy students and pharmacy residents. She is also a clinical Assistant Professor in the School of Nursing where she teaches oncology obtaminacology in the purse practitioner program.	
7	
8	_
OUR PRESENTER	
PRE	
8	
]
Preceptor to pharmacy students and pharmacy residents.	
OUR PRESENTER	
PRES	
•	

	5 .	Preceptor to pharmacy students and pharmacy residents.
OUR PRESENTER	4.	Clinical Assistant Professor in the School of Nursing where she teaches oncology pharmacology in the nurse practitioner program.
10		

5 Preceptor to pharmacy students and pharmacy residents.

4. Clinical Assistant Professor in the School of Nursing where she teaches oncology pharmacology in the nurse practitioner program.

3. Adjuvant Faculty at Eshelman School of Pharmacy where she teaches breast cancer treatment to pharmacy students.

11

Preceptor to pharmacy students and pharmacy residents. Clinical Assistant Professor in the School of Nursing where she teaches oncology pharmacology in the nurse practitioner program. Adjuvant Faculty at Eshelman School of Pharmacy where she teaches breast cancer treatment to pharmacy students. In 2014, joined Lineberger's outpatient breast oncology team.

i		
	5 .	Preceptor to pharmacy students and pharmacy residents.
i		
TER	4.	Clinical Assistant Professor in the School of Nursing where she teaches oncology pharmacology in the nurse practitioner program.
	•	Adiment Ferrity at February Cahaal of Dharman
PRESENTER	3.	Adjuvant Faculty at Eshelman School of Pharmacy where she teaches breast cancer treatment to pharmacy students.
~		In 2014, initial Linebarger's outpetient breest
	2.	In 2014, joined Lineberger's outpatient breast oncology team.
	4	Since 2008, Clinical Pharmacist Practitioner
	ı.	at the North Carolina Basnight Cancer Hospital.
		g



14

This activity has been planned and implemented under the sole supervision of the Course Director, William A. Wood, xo, xn+, in association with the UNC Office of Continuing Professional Development (CPD). The course companies as defined by the ACCME.

The University of North Carolina at Chapel Hill is accredited with distinction as a provider of nursing continuing professional development by the American Nurses Credenialing Center's Commission on Accreditation. A potential conflict of interest sociars when an individual has an opportunity to affect educational content about health-care products or services of a social content of the conflict of t





17

Oral Chemotherapy Objectives • Describe the factors involved in safe prescribing practices of oral chemotherapy • Review how to appropriately monitor patients while on oral chemotherapy • Discuss the safe handling, storage and disposal of oral chemotherapy

Oral chemotherapy

- Oral chemotherapy includes medications taken by mouth that are cytotoxic or target proteins, pathways or receptors that are involved with cancer
- Use of oral chemotherapy has been steadily increasing
 18 novel oral chemotherapies have been approved from 2020 to
 December 2022
- Estimated that 25-30% of chemotherapy medications in the development pipeline will be oral

ugs at FDA: CDER's New Molecular Entities and New Therapeutic Biological Products. https://www.fda.gov/dnogdevelopment-approval-process-drugs/new-drugs-fda-cdersnew.moleculs.neotities.and.new.fherapeutic biological products. Acressed Therember 8, 2022

19

Shifts the care of patients from the provider to the patient

Benefits

Concerns

- Patient convenience
- Adherence
 Safety
- Patient empowerment
- Proper storage/handling
- Efficacious
- Monitoring
- · Reduced toxicity?
- Cost

Weingart SN, Toro J, Spencer J, et al: Medication errors involving oral chemotherapy. Cancer 216:2455-2464, 202:
Weingart SN, et al: NCCNTask Force: Oral Chemotherapy. J Natl Comp Canc Netw 2008:6 Suppl 3:51-14.

20


Wide scope of use CML Imatinib, nilotinib, dasatinib, bosvitnib Metastatic RCC Ph. AML Imatinib, pazopanib, cabozantinib, axitnib BRAF-mutant melanoma ALK-mutant non-small cell lung cance Crizotinib, alectinib, dastrafenib/trametinib Multiple myeloma EGFR-mutant NSCLC Advanced prostate Cancer Advanced breast cancer Advanced breast cancer Advanced colon cancer Advanced colon cancer Advanced colon cancer Capecitabine, regorafenib, trifluridine + tipiracil

Varied mechanisms of action			
	Cytotoxic	• E.g. Capecitabine	
	Targeted	• E.g. Imatinib	
	Endocrine	• E.g. Tamoxifen	
	Other	• E.g. Lenalidomide	



23







Prescribing Oral chemotherapy prescriptions should include: Patient name and demographics Dose calculation, if applicable Schedule Administration instructions Quantity Refills Phases Septiment of the se


26

Patient education Patients should be provided education prior to initiation of oral chemotherapy Best to use standardized teaching materials Should be suitable for patient's health literacy and language Can clarify patient misconceptions Will need reinforcement throughout treatment period





29



Education should be provided verbally and in written form Patient education family members, caregivers or others Patients should be encouraged to write down questions and answers

31

Patient education resources

- Website resources that provide handouts specifically for patients

- Oncolink (https://www.oncolink.org/)
 Chemocare (https://chemocare.com/)
 Oral Chemotherapy Education (https://www.oralchemoedsheets.com/)
- Package insert
- Tertiary drug information databases
- Lexi-Comp, UpToDate®, Facts and Comparisons, Micromedex
- www.drugs.com/monograph
 Free website that obtains drug information from multiple drug information databases including Micromedex and American Society of Health-System Pharmacists

32

Drug interactions

- Oral chemotherapy can interact with other medications,
- supplements, or foods
- Can result in increased toxicities or reduced efficacy
- Interactions may occur by multiple mechanisms
- Important to assess for interactions at each visit
- Information on interactions and their management may be

Segal EM et al. Oral chemotherapy food and drug interactions: A comprehensive review of the literature. J Oncol Pract 2015;10: e255-268 Rogala BD et al. Oral anticancer therapy: Management of drug interactions. J Oncol Pract 2019;15;81-90

Pharmacokinetic drug interactions

- · Effect the absorption, distribution, metabolism or excretion of another
- CYP 3A4 is the most common enzyme that metabolizes oral chemotherapy • Drugs that inhibit or induce CYP 3A4
- can effect concentrations of drugs that are substrates of this enzyme

EXAMPLES

- Phenytoin (Dilantin®)
- Strong 3A4 inducer
- Can significantly decrease the concentrations of medications that are major substrates of CYP 3A4
- Nirmatrelvir/ritonavir (Paxlovid®)
- Strong 3A4 inhibitor
- Can significantly increase the concentration of major substrates of CYP 3A4

Segal EM et al. Oral chemotherapy food and drug interactions: A comprehensive review of the literature. J Oncol Pract 2015;10: e255-268 Rogala BD et al. Oral anticancer therapy: Management of drug interactions. J Oncol Pract 2019;15:81-90

34

EXAMPLE

Pharmacodynamic drug interactions

Lead to an alteration of pharmacologic effect, which may be additive, synergistic, or antagonistic

QTc prolongation

- Oral chemotherapy agents may prolong QTc and the addition of other drugs that have this side effect result in an additive risk
- Ribociclib (Kisqali ®) can cause QTc prolongation. Addition of ondansetron (Zofran ®) or citalopram (Celexa ®) can increase the risk of this adverse effect




35


EXAMPLES

Acid suppression therapy interactions

PPIs, H2 receptor antagonists, and antacids can effect the absorption of some oral chemotherapy

- Proton pump inhibitors
- Due to decreased absorption, PPIs are contraindicated with dasatinib, erlotinib, neratinib and pazopanib
- Histamine H2 receptor antagonists
 Erlotinib, and nilotinib can be taken 10 hours after or 2 hours before taking a histamine H2-receptor
- Antacids
 Over-the-counter antacids should be administered hours before or after the administration of acalabrutinil, dasatinily, ertotinil, geftinili, bosunilin, initiniil, and ponatinib, and 3 hours before neratinib






38


Drug interaction resources


- Manufacturer package insert
- Tertiary drug information databases
- Lexi-Comp, UpToDate®, Facts and Comparisons, Micromedex
 www.drugs.com/drug_interactions.html
- Herb information
- Natural Medicines, available through institutional library
 www.naturalmedicines.therapeuticresearch.com
- Memorial Sloan Kettering Cancer Center website About Herbs, Botanicals & Other Products
- Now.mskc.corg/cancer-care/diagnosis-treatment/symptom-management/integrative-medicine/herbs
 National Center for Complimentary and Integrative Health
- www.nccih.nih.gov







41









44









47






Resources to help with cost Charity Grants Patient Access Network Foundation www.panfoundation.org Healthwell https://www.healthwellfoundation.org/ Patient Advocate Foundation https://conavs.org/ NeedyMeds www.needymeds.org 10fUS 1ofus.org Leukemia Lymphoma Society www.lls.org Social Security Compassionate Allowance www.socialsecurity.gov/compassionateallowances/ Pharmacists


50







53



Potential labs/tests to monitor



- CBC/diff
- LFTs
- SCr
- Electrolytes
- · Potassium/magnesium Lipid panel and glucose
- · Lipase/amylase
- Uric acid

- CPK • TSH
- Pregnancy
- ECG
- ECHO
- · Blood pressure

55

Where to find required lab testing and intervals

Tertiary drug information databases

 Lexi-Comp UpToDate® · Facts and Comparisons Micromedex • Drugs.com DailyMed

Package insert

Imatinib (Gleevec®)

- WARNINGS AND PRECAUTION

56

Example: Imatinib

- Pretreatment Screening
 Complete blood cell count (CBC)
 Baseline liver and renal function
 Consider performing an echocardiogram and determining serum troponin concentrations in patients with hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukemia (CEL), and in patients with hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukemia (CEL), and in patients with myelodysplastic/myeloproliferative diseases or aggressive systemic castocytosic sasociated with high eosinophil levels. It results of the echocardiogram or serum troponin concentrations are abnormal, consider prophylactic use of systemic confocations.

 Verify pregnancy status in females of reproductive potential

Patient Monitoring

- **Monitor for signs or symptoms of fluid retention (e.g., weight gain) regularly during therapy

 Monitor GBC weekly for the first month of therapy, every other week during the second month, and periodically
 (e.g., every 2–3 months) thereafter as clinically indicated

 Carefully monitor patients with critical disease or risk factors for cardiac disease for signs and symptoms of

 cardiac toxicity or renal failure

- cardiac toxicity or renal failure

 Cardiul's monitor patients with a history of renal failure for signs and symptoms of cardiac toxicity or renal failure


 Monitor liver function monthly or as clinically indicated during therapy

 Monitor renal function during therapy


 Monitor serum TSH concentrations in patients receiving levothyroxine replacement therapy following thyroidectomy
 in pediatric patients, monitor bone growth and development


 Monitor patients with high tumor burden or those with a high proliferative rate for TL Satps, livewed-ups, continuous pashfirmatish.tn


Monitoring CML TKIs							
	Imatinib	Nilotinib	Dasatinib	Bosutinib	Ponatinib	Asciminib	
СВС	X (weekly x1 mo→biweekly x1 mo→periodically)	X (biweekly x2mo → monthly)	X (weekly x12 weeks → q3 months)	X (weekly xamo→ monthly)	X (biweekly x3mo → monthly)	X (biweekly x3mo → monthly)	
LFTs	X (baseline→ monthly)	X (baseline→ monthly)	X (baseline → periodically)	X (monthly x3→ periodically)	X (baseline → monthly)		
SCr	X (baseline→ periodically)			X (baseline → periodically)			
Electrolytes	X (baseline→ periodically)	X (baseline→ periodically)	X (baseline → periodically)		X (baseline → periodically)		
Lipid profile & glucose		X (baseline → Periodically)					
Lipase & Amylase		X (baseline → monthly)			X (biweekly x2mo → monthly)	X (baseline → monthly)	
Uric Acid		X (baseline)			X (baseline)		
ECG		X (baseline → periodically)	X (baseline→ q3mo x2)				
Pregnancy	Х	X	X	X	X	Х	




59










62







65





Safe handling Recommendations for health care providers

- Receive and maintain training on safe practices if the practitioner will come into contact with oral chemotherapy
- Follow institutional policies
- Follow guidelines related to oral chemotherapy

Goodin, S et al. Safe handling of Oral Chemtherapeutic agents in Clinical Practice: Recommendations from an International Pharmacy panel. JOP 2011;7(1): 712.

68



- Understand handling and administration instructions
- Tablets/capsules should not be crushed, chewed, or cut
- Use gloves or tip tablets into a disposable medicine cup
- Wash hands thoroughly
- If able to place in a pill box, do not share the pill box with others
- \bullet Double flush or use a separate bathroom, if able

Goodin, S et al. Safe handling of Oral Chemterapeutic agents in Clinical Practice: Recommendations from an International Pharmacy panel. JOP 2021;7(2): 73

Safe storage



- Store in a safe place away from children and pets
- Store medications away from sunlight, humid bathrooms, or other areas that may have fluctuations in temperature
- Read labels some oral chemotherapy must be stored in original container (e.g. regorafenib) or refrigerated (e.g. oral etoposide, trametinib)
- If traveling, store in original, pharmacy-labelled container

70

Safe disposal

- Don't discard unused tablets/capsules down the toilet
- Dispose in local medication disposal sites or drop boxes
- May mix whole tablets with something unpalatable (e.g. coffee grounds) and place in a container with lid or double sealable storage bags prior to placing in regular trash
- Celgene provides packaging so patients may return unused lenalidomide, pomalidomide, and thalidomide

71

Oral chemotherapy resources

Hematology/Oncology Pharmacy Association (HOPA)
 Oral chemotherapy resources
 www.hoparx.org/images/hopa/resource-library/guidelinesstandards/Oral Chemo Resources Web Outline.pdf.

- Oncology Nursing Society
 Oral Anticancer Medication Toolkit
- Drug.com
 Drug monographs
 www.drugs.com/


- Pharmacists






74









77







80

